

Fuzzy Bags and Wilson Lines

Robert D. PISARSKI

*Department of Physics
Brookhaven National Laboratory
Upton, NY 11973 USA*

I start with an elementary observation about the pressure in the deconfined phase of a SU(3) gauge theory without quarks. This suggests a “fuzzy” bag model for the analogous pressure in QCD, with dynamical quarks. I then sketch how the deconfined phase might be described using an effective theory of Wilson lines. To leading order in weak coupling, the effective electric field appears in a form familiar from the lattice theory of Banks and Ukawa.

§1. Fuzzy Bags

The spectacular success of the heavy ion programs at the SPS and RHIC justifies a careful analysis of the phase transition(s) of QCD at nonzero temperature. In this Proceeding I summarize some recent work of mine,¹⁾ hopefully in a more comprehensible fashion.

Any fundamental understanding of these phase transitions rests upon the bedrock provided by numerical simulations of lattice QCD. In this section I begin by looking at old data²⁾ in a new way. This was mentioned in a footnote, Ref. 37 of Ref. 1. While the data is for a pure gauge theory, it shows that except very near T_c , the critical temperature, there is an exceedingly simple form for the pressure. This immediately suggests an ansatz applicable to QCD, and which might be of use for phenomenology.




Fig. 1. Lattice results²⁾ for the pure SU(3) gauge theory: to the left, $(e-3p)/T^4$; to the right, the same quantity times T^2 .

Ten years ago, a group at Bielefeld²⁾ computed the thermodynamic properties of a SU(3) gauge theory, close to the continuum limit. In equilibrium, while all thermodynamic quantities follow from the pressure, it is convenient to plot what is

usually called the “interaction measure”, $(e - 3p)/T^4$, where e is the energy density, p is the pressure, and T the temperature. This is plotted in the left panel in Fig. 1. The interaction measure is the trace of the energy momentum tensor, divided by T^4 , and so vanishes if the theory is conformally symmetric. A pedestrian way of seeing this is to note that $e - 3p = T(dp/dT) - 4p$; if the theory is conformally invariant, the pressure is just a pure number times T^4 , for which $e - 3p$ vanishes.

Thus the interaction measure is a dimensionless number which quantifies the deviation from conformal symmetry. Because the conformal anomaly is proportional to the β -function, this also measures the deviation from ideality. As can be seen from the left panel, the interaction measure is very small below T_c , rises steeply around T_c , with a sharp peak at $T_{\max} \approx 1.1 T_c$. Above $T_{\max}$ it trails off relatively slowly. By $T_{\text{pert}} \approx 4.0 T_c$, its value is, within a factor of two, equal to that expected from perturbation theory,¹⁾ where the interaction measure begins as $\sim \alpha_s^2$.

My concern is with the fall off between $T_{\max}$ and T_{pert} . In the right panel of Fig. 1, I take $(e - 3p)/T^4$, and simply multiply times T^2 . As can be seen, $(e - 3p)/T^2$ is essentially constant. For the temperatures shown, then, the pressure is a sum of just *two* pieces: an ideal term, $\sim T^4$, and a new, non-ideal term, $\sim T^2$.

If one ignores the overall normalization, the relative normalization of the two terms can be computed without further ado. In an asymptotically free theory, at high temperature the pressure approaches that of an ideal gas, so at any temperature it is natural to compare the pressure to that of the appropriate ideal gas. In the pure glue theory, $T_c \sim 270$ MeV is much smaller than the lightest glueball mass, ~ 1.5 GeV, and so, relative to the ideal gas, the confined pressure is very small.^{2),3),4)} Thus the pressure (nearly) vanishes at T_c . This gives

$$p_{\text{pure glue}}(T) \approx f_{\text{pert}} (T^4 - T_c^2 T^2) \quad , \quad 1.1 T_c \leq T \leq 4.0 T_c \quad , \quad (1.1)$$

where f_{pert} is a constant.

What of other numbers of colors, N_c ? The deconfining phase transition is of second order³⁾ for $N_c = 2$, and of first order^{2),4)} when $N_c \geq 3$. As N_c increases, the transition becomes more strongly first order,⁴⁾ with a latent heat $\sim N_c^2$. For three colors,²⁾ the transition is weakly first order.

The change in the order of the transition with N_c affects the interaction measure, but not dramatically so.⁴⁾ Since the pressure is continuous at T_c , and as the confined phase has negligible pressure, then whatever the order, the pressure is almost zero at T_c , $p(T_c) \approx 0$. In contrast, the energy is sensitive to the order: while $e(T_c^-) \approx 0$, just above the transition the energy vanishes for a second order transition, and is nonzero for a first order transition.

For a first order transition, then, at T_c the interaction measure $\approx e(T_c^+)/T_c^4$. For the strongly first order transition with four colors,⁴⁾ it is not surprising to find that the maximum in the interaction measure is at $T_c = T_{\max}$, and that it falls off after that.

For a second order transition, the energy is continuous at T_c , so $e(T_c^+) \approx 0$, and the interaction measure is nearly zero at T_c . For the deconfining transition with two colors,³⁾ the interaction measure increases from near zero at T_c , has a sharp maximum at $T_{\max} \approx 1.15 T_c$, and then falls off after that. Thus for two colors,³⁾ the

peak in the interaction measure can be viewed as a remnant of that for an infinite number of colors.⁴⁾ Since the transition is nearly second order for three colors,²⁾ $e(T_c^+)$ is small, and the interaction measure looks like that of two colors; the peak in the interaction measure moves closer to T_c , to $T_{\max} \approx 1.1 T_c$.

The formula in (1.1) only applies above the maximum in the interaction measure, so we should only compare different N_c at $T > T_{\max}$. For $N_c = 4$, this is for all $T \geq T_{\max} = T_c$. This appears to be supported by lattice simulations.⁴⁾ To be fair, the data for $N_c = 8$ does not, but perhaps lattice discretization errors are larger there.⁴⁾ For two colors, one compares for $T > T_{\max} = 1.15 T_c$: the data of Ref. 3 appears to violate (1.1) by a large amount, $\sim 50\%$. However, it is not clear that these simulations are close to the continuum limit, and so new simulations would be most welcome.

Before describing how one might extend (1.1) to other temperatures, to $T < T_{\max}$ and $T > T_{\text{pert}}$, I skip ahead directly to the case of dynamical quarks and three colors. For three flavors or less,⁵⁾ while the “transition” often becomes a crossover, an approximate “ T_c ” can still be defined. Whatever the order, though, the interaction measure behaves similarly. Relative to an ideal gas of quarks and gluons, there is a small but significant pressure below T_c , and a maximum in the interaction measure at a temperature $T_{\max}$, which is above T_c . The surprise about (1.1) is that the leading correction to the ideal gas term is $\sim T^2$, and not $\sim T^3$. I speculate that this is generic: that for $T_{\max} < T < T_{\text{pert}}$, the pressure is a series in powers of $1/T^2$ times the ideal T^4 term. I call this a “fuzzy” bag model for the pressure:

$$p_{\text{QCD}}(T) \approx f_{\text{pert}} T^4 - B_{\text{fuzzy}} T^2 - B_{\text{MIT}} + \dots, \quad T_{\max} \leq T \leq T_{\text{pert}} \quad (1.2)$$

The upper bound, T_{pert} , denotes the point at which perturbative contributions to the interaction measure are of the same order as that found from the lattice; it is a few times T_c , something like $\approx 4T_c$.

In (1.1), f_{pert} is dimensionless, B_{MIT} is the usual MIT bag constant,⁶⁾ with dimensions of $(\text{mass})^4$, and B_{fuzzy} is a fuzzy bag constant, with dimensions of $(\text{mass})^2$. For the pure glue theory, from (1.1) $B_{\text{fuzzy}} = f_{\text{pert}} T_c^2$, and $B_{\text{MIT}} \ll B_{\text{fuzzy}}$.

Recent lattice simulations⁵⁾ appear to support (1.2). Note that with a fuzzy bag constant, the sign of the MIT bag constant is not guaranteed, but these simulations find that B_{MIT} is positive,⁵⁾ as in the original MIT bag model.⁶⁾

With a little work, it should be possible to generalize (1.2) to the entire deconfined phase. Above T_{pert} , we can ignore the non-ideal terms, and include only f_{pert} , now considered as a function of T . This is given by resummations of weak coupling perturbation theory at nonzero temperature.^{7),8)} While all resummations fail at temperatures below T_{pert} , in the present view this is just because of the *non*-ideal terms in the pressure. This suggests that for all $T \geq T_c$, perturbative resummations contribute only to $f_{\text{pert}}(T)$. Analysis shows that $f_{\text{pert}}(T_{\text{pert}})$ is about 90% of the ideal gas value.^{7),8)} If applied just to $f_{\text{pert}}(T)$, it seems very possible that perturbative resummation might work all of the way down to the critical temperature.⁷⁾ With present day techniques, lattice simulations could test this in a precise manner in the pure glue theory.

This suggests a heuristic analogy, to the operator product expansion for two

gauge invariant operators at short distances. Free field theory dominates as the distance $x \rightarrow 0$, like some power of x . Perturbative corrections enter with the same power, times a series in $1/\log(x)$, *etc.* Non-perturbative effects involve the expectation values of new operators, multiplied by the appropriate powers of x^2 . For the pressure, perturbative terms correct the ideal T^4 term, as a series in $1/\log(T)$, *etc.*, while non-perturbative effects generate non-ideal terms, as a series in $\sim 1/T^2$ times the ideal term.

What about below $T_{\max}$? I assume that a hadron resonance gas provides a reasonable approximation not just about $T = 0$, but all of the way to T_c . This leaves $T_c \leq T \leq T_{\max}$. For three colors in the pure glue theory, this is a nearly critical regime, dominated by a light excitation for the triplet Polyakov loop. Maybe even with dynamical quarks, this region is dominated by a light triplet loop and its interactions with quarks.⁹⁾

Why bother? For a pure glue theory, simulations relatively close to the continuum limit have been possible for some time.²⁾ With dynamical quarks, though, and in particular for the light quarks present in QCD, present day simulations are not close to the continuum limit. Thus (1.2) could be used by approximate models. For example, most hydrodynamic models¹⁰⁾ use a MIT bag model for the pressure. This approaches ideality much faster than a fuzzy bag model. Since hydrodynamics only requires the energy as a function of pressure, perhaps non-ideal corrections don't really matter that much, but this should be demonstrated by explicit computation.

Terming (1.2) a “fuzzy” bag is not sheer whimsy. In the MIT bag model the surface of the bag is infinitely thin and fixed.⁶⁾ While the interface surely has nonzero width, it is difficult to know how to model this. For instance, if the surface of the bag were thin and floppy, then there would be many light excitations, in which the surface of the bag flops around, and the quarks remain essentially fixed.¹¹⁾ There are no signs of such a multiplicity of states in the hadronic mass spectrum.

The essential moral of the non-ideal terms in the pressure, (1.1) and (1.2), is that the transition from a confined, to a nearly perturbative phase, is not abrupt, as in a MIT bag model, but gradual. This suggests that the boundary of the bag isn't thin, but thick. A thick bag is unlikely to flop around, since the entire width needs to participate. Thus light surface modes shouldn't be a problem.

Having said this, it is not clear how to develop a more realistic bag, with a thick boundary. The thickness probably doesn't affect the mass spectrum of ordinary hadrons greatly. The analogy is still suggestive. In some loose sense nonzero temperature probes inverse distances in the QCD vacuum, $T \sim 1/R$: the perturbative vacuum emerges as $T \rightarrow \infty$, or $R \rightarrow 0$; the confined vacuum, for $T \rightarrow 0$, or $R \rightarrow \infty$. The width of the bag emerges over distances $R \approx 1/T_{\text{pert}} \rightarrow 1/T_c$; in physical units, for $\approx 1/4 \text{ fm} \rightarrow 1 \text{ fm}$. This is the right scale to probe the transition from going inside, to outside, the bag.

§2. Wilson lines and their electric field

The above begs the question: what is the origin of the non-ideal terms in the pressure of the deconfined phase? I next turn to a possible explanation in terms of

Wilson lines.¹⁾ I omit all details, and many references, to concentrate on a broad and qualitative description.

In resummations of perturbation theory,^{7),8)} the gluon degrees of freedom remain the time-like, A_0 , and space-like, A_i , components of the vector potential. To proceed further, consider a straight, thermal Wilson line:

$$L(x) = P e^{ig \int_0^{1/T} A_0(x, \tau) d\tau} . \quad (2.1)$$

$\tau : 0 \rightarrow 1/T$ is the imaginary time, x is the spatial coordinate, and g the gauge coupling constant. If A_0 is in the fundamental representation, L transforms as an adjoint matrix: under strictly periodic gauge transformations, $\mathcal{U}(x, 0) = \mathcal{U}(x, 1/T) = \mathcal{U}(x)$, $L(x) \rightarrow \mathcal{U}^\dagger(x) L(x) \mathcal{U}(x)$.

The trace of the Wilson line is the Polyakov loop, and is invariant under local gauge transformations. The motivation for considering an effective theory of the Wilson line rests upon measurements of the renormalized Polyakov loop. In a perturbative regime, fluctuations in A_0 should be small, and so suitably normalized, this expectation value should be near one. In contrast, if fluctuations in A_0 are large, the expectation of the Polyakov loop is not near one. Lattice simulations in a SU(3) gauge theory, with or without quarks, show the (renormalized, triplet) loop is near one for $T > T_{\text{pert}}$, and less than one for $T_c < T_{\text{pert}}$.

One's first guess might be that the theory is driven into a regime of strong coupling. Consider, however, the ‘‘Helsinki’’ program of resummation.⁸⁾ Originally proposed by Braaten and Nieto, computations to four loop order were done by Kajantie, Laine, Rummukainen, and Schröder. The final steps are being completed by Laine, Schröder *et al.*⁸⁾ They find that the effective coupling runs with a mass scale $\sim 2\pi T$, so that even down to $T_c \sim 175$ MeV, the QCD coupling is $\alpha_s^{\text{eff}}(1.6 \text{ GeV}) \sim 0.28$. This value is *not* that large.⁸⁾

This suggests the perturbative construction of an effective theory in three dimensions, valid over distances $> 1/T$. Since the renormalized Polyakov loop is not near unity, A_0 is replaced by Wilson lines, coupled as always to the A_i .

When I first suggested this there were several problems.⁹⁾ The first is how one to match the effective theory, at large A_0 , to quantities which are computable perturbatively. In the Helsinki program, this is done by computing the positions of poles in propagators, *etc.* This is fine at small A_0 , but doesn't probe large A_0 .

Interfaces¹²⁾ can be used to probe large A_0 . These are most familiar in a pure gauge theory, such as SU(N). A SU(N) gauge group has a global center symmetry of Z(N), so that in the deconfined phase, there are N degenerate vacua: the usual perturbative vacua, $L = 1_N$, and Z(N) transforms thereof, such as $L = \exp(2\pi i/N) 1_N$.

An interface interpolates between these degenerate vacua. One takes a box which is long in one spatial direction, say $z : 0 \rightarrow z_f$. At one end of the box, one takes one vacua, $L(0) = 1_N$; at the other end, an inequivalent Z(N) state, $L(z_f) = \exp(2\pi i/N) 1_N$. These boundary conditions force the formation of an interface along z , which tunnels between the degenerate vacua. While the ends of the box represent perturbative vacua, in between one probes large $A_0 \sim T/g$, as illustrated by (2.3). The amplitude for tunneling can be computed semiclassically, and gives

an interface tension $\sim T^2/\sqrt{g^2}$. The width of the interface is proportional to the inverse Debye mass, $\sim 1/(\sqrt{g^2}T)$, so that a derivative expansion can be used. To date, computations have been carried out to $\sim (\sqrt{g^2})^{3/2}$ times the result at leading order by Giovannangeli and Korthals Altes.¹²⁾

Such interfaces appear to be special to theories with a center symmetry, and so useless for QCD, where it is violated by the presence of dynamical quarks. In this case, however, there are other interfaces which can be used. Consider the gauge transformation

$$\mathcal{U}_c(\tau) = e^{2\pi i \tau T t_N/N} \quad , \quad t_N = \begin{pmatrix} 1_{N-1} & 0 \\ 0 & -(N-1) \end{pmatrix} . \quad (2.2)$$

This is only periodic up to a $Z(N)$ rotation, $\mathcal{U}_c(1/T) = \exp(2\pi i/N) \mathcal{U}_c(0)$, which is allowed in the absence of quarks. If one acts with $\mathcal{U}_c$ on one end of the box, but not the other, then a $Z(N)$ interface forms, because this isn't a pure gauge transformation in between the two ends.

Now instead of (2.2), consider the N^{th} power thereof, $\mathcal{U}_c^N$. This is a strictly periodic gauge transformation, and thus is allowed, independent of whatever matter fields may be present. If we act with $\mathcal{U}_c^N$ on one end of a box, but not the other, what I term a $U(1)$ interface is generated: while $\langle L \rangle = 1_N$ at both ends, in between A_0 winds around in a topologically nontrivial fashion.

To compute the properties of an interface, one needs the effective potential for constant A_0 , which is first generated at one loop order. For a effective theory of Wilson lines, it is obvious to turn an effective potential of A_0 into one for L . What stymied me⁹⁾ is what one does at *zeroth* order: how does one write the effective electric field in terms of Wilson lines?

The crucial clue was provided by what appeared to be an abstruse computation. At one and two loop order in a $SU(N)$ gauge theory, explicit calculation shows that the effective potential, computed in the presence of a large, background field for constant A_0 — and $A_i = 0$ — is invariant under the $Z(N)$ center symmetry.¹²⁾ This is unremarkable: there is no anomaly to prevent the quantum theory from respecting the center symmetry of the classical theory. Diakonov and Oswald¹³⁾ then computed in the presence of background fields in which both A_0 and A_i were nonzero, allowing A_0 to be large. Assuming that the effective electric field is $D_i A_0$, they found that the center symmetry appeared to be violated at one loop order. An error in computation seems unlikely, given that their results agree with those of Megias, Ruiz, and Salcedo,¹⁴⁾ who computed in the limit of small A_0 and A_i .

I suggest that the problem lies in assuming that the effective electric field is $D_i A_0$, and arises even at tree level. Under the large gauge transformation of (2.2),

$$A_0^{diag} \rightarrow A_0^{diag} + \frac{2\pi T}{gN} t_N \quad , \quad A_i \rightarrow \mathcal{U}_c^\dagger A_i \mathcal{U}_c . \quad (2.3)$$

Hence diagonal elements of A_0 are shifted by a large but constant amount, $\sim T t_N/g$, while off-diagonal elements of A_0 and A_i undergo time dependent rotations.

In four dimensions, the original electric field is $D_i A_0 - \partial_0 A_i$. The first term, $D_i A_0 = \partial_i A_0 - ig[A_i, A_0]$, changes if $[A_i, t_N] \neq 0$, which always happens for some

(off-diagonal) components of A_i . It is easy to see, however, that the time dependent rotation of A_i in the second term, $-\partial_0 A_i$, generates a similar commutator, $-[A_i, t_N]$, which exactly cancels against the first term. This just reflects the fact that the original electric field transforms homogeneously under arbitrary gauge transformations.

In the effective theory, the simplest guess for the effective electric field is to drop time derivatives, and take it to be $D_i A_0$. While valid at small A_0 ,^{(7),(8)} this can't be right at large A_0 . The above argument shows that $D_i A_0$ is not invariant under the large gauge transformations, $\mathcal{U}_c$, which enforce the center symmetry.⁽¹³⁾ Similarly, if we take $\mathcal{U}_c^N$, the effective theory is not even invariant under large, but strictly periodic, gauge transformations.

The resolution is to construct the effective electric field from Wilson lines. These transform like a phase under $\mathcal{U}_c$, and are invariant under $\mathcal{U}_c^N$. To leading order in weak coupling, the effective electric field is

$$E_i^{\text{eff}}(x) = \frac{T}{ig} L^\dagger(x) D_i L(x). \quad (2.4)$$

This is shown by demonstrating that the interface tensions, either $Z(N)$ or $U(1)$, agree between the effective and original theories. Eq. (2.4) does change beyond leading order, and is multiplied by other gauge invariant terms, such as $|\text{tr } L|^2$, $|\text{tr } L^2|^2$, etc.; with coefficients which begin at $\sim g^2$.

It is well known that the mapping between the fields in an effective theory, and the original theory, is indirect. Usually, however, one only sees this at next to leading order in some expansion, so the differences are small. In the present instance, because one is constructing an effective theory for large A_0 , it arises even at zeroth order. The problems found before⁽¹³⁾ are presumably solved by matching to an effective theory constructed from the effective electric field, and not functions of $D_i A_0$.

To leading order, the effective Lagrangian is that of a gauged, principal chiral field:

$$\mathcal{L}_{\text{classical}}^{\text{eff}}(A_i, L) = \frac{1}{2} \text{tr } G_{ij}^2 + \frac{T^2}{g^2} \text{tr } |L^\dagger D_i L|^2. \quad (2.5)$$

This is nonrenormalizable, but this shouldn't be a problem, since the effective theory is only valid over distances $> 1/T$. A related linear model, constructed to be renormalizable, has been analyzed by Vuorinen and Yaffe.⁽¹⁵⁾

On the lattice, the Lagrangian of (2.5) was first written down by Banks and Ukawa,⁽¹⁶⁾ as the simplest kinetic term for an adjoint field, L . It is not obvious that (2.5) applies in the continuum, even at leading order in weak coupling. It should be possible to construct effective Lagrangians to the same order as interfaces, in the original theory, have been computed.⁽¹²⁾

One can show that the four dimensional instanton number equals the winding number of the Wilson line.⁽¹⁾ Dai and Nair⁽¹⁷⁾ showed that non-abelian hydrodynamics has color Skyrmions. This suggests that the effective model might have electric Skyrmions:⁽¹⁷⁾ solutions stabilized by a nonzero winding number for L , and yet which are not instantons, since they only carry electric, and not magnetic, fields.

How does the deconfining transition arise? For the potential for L computed perturbatively, order by order the deconfined vacuum is always proportional to the

unit matrix, $\langle L \rangle \sim 1_N$. This could be possible all of the way to T_c — but then one would not expect non-ideal terms in the pressure. To destabilize the perturbative vacuum, it is necessary to add, by hand, a non-perturbative mass term for the Wilson line. The simplest example is $\sim T^2 B_f |\text{tr } L|^2$. The notation B_f , and the T^2 , is motivated by the previous section; adding such a mass term is standard in a Landau-Ginzburg type of analysis.

With such a term, $\langle L \rangle \neq 1$. In Ref. 1, I argued that at infinite N_c , the confined vacuum is characterized by complete eigenvalue repulsion; also, that lattice results suggest that this is approximately true at small N_c . The appearance of eigenvalue repulsion is clear when the spatial volume is small, as when the theory lives on a very small sphere. At infinite N_c , Aharony *et. al.*¹⁸⁾ showed that the effective theory only involves a constant mode, and reduces to a random matrix model for that mode. As typical of random matrix models, eigenvalue repulsion arises from the Vandermonde determinant in the measure of the group integral.

In infinite volume, instead of one random matrix, there is a field theory of (not quite) random matrices. Such field theories have been studied little. The interesting, and gauge invariant, question is how confinement arises from the *dynamical* generation of eigenvalue repulsion.

§3. Acknowledgements

This research was supported by D.O.E. grant DE-AC02-98CH10886, and in part by the Alexander von Humboldt Foundation. I thank Prof. T. Kunihiro, and the other organizers of YKIS 2006, for their invitation and warm hospitality.

References

- 1) R. D. Pisarski, to appear in Phys. Rev. D [[arXiv:hep-ph/0608242](#)].
- 2) G. Boyd, J. Engels, F. Karsch, E. Laermann, C. Legeland, M. Lutgemeier, and B. Petersson, Nucl. Phys. B **469** (1996), 419 [[arXiv:hep-lat/9602007](#)].
- 3) J. Engels, J. Fingberg, K. Redlich, H. Satz, and M. Weber, Zeit. fur Phys. **42** (1989), 341.
- 4) B. Bringoltz and M. Teper, Phys. Lett. B **628** (113), 2005 [[arXiv:hep-lat/0506034](#)]; B. Bringoltz, private communication.
- 5) M. Cheng *et al.*, [[arXiv:hep-lat/0612001](#)]; F. Karsch, private communication.
- 6) A. Chodos, R. L. Jaffe, K. Johnson, C. B. Thorn, and V. F. Weisskopf, Phys. Rev. D **9** (1974), 3471.
- 7) J. O. Andersen and M. Strickland, Ann. Phys. **317** (2005), 281 [[arXiv:hep-ph/0404164](#)]; U. Kraemmer and A. Rebhan, Rept. Prog. Phys. **67** (2004), 351 [[arXiv:hep-ph/0310337](#)]; A. Rebhan, private communication.
- 8) M. Laine and Y. Schröder, J. High Energy Phys. **0503** (2005), 067 [[arXiv:hep-ph/0503061](#)]; F. Di Renzo, M. Laine, V. Miccio, Y. Schröder and C. Torrero, J. High Energy Phys. **0607** (2006), 026 [[arXiv:hep-ph/0605042](#)].
- 9) R. D. Pisarski, Phys. Rev. D **62** (2000), 111501(R) [[arXiv:hep-ph/0006205](#)].
- 10) P. Huovinen and P. V. Ruuskanen, [[arXiv:nucl-th/0605008](#)].
- 11) R. L. Jaffe, private communication at YKIS 2006.
- 12) P. Giovannangeli and C. P. Korthals Altes, Nucl. Phys. B **721** (2005), 1 [[arXiv:hep-ph/0212298](#)]; **721** (2005), 25 [[arXiv:hep-ph/0412322](#)].
- 13) D. Diakonov and M. Oswald, Phys. Rev. D **68** (2003), 025012 [[arXiv:hep-ph/0303129](#)]; **70** (2004), 105016 [[arXiv:hep-ph/0403108](#)].
- 14) E. Megias, E. Ruiz Arriola, and L. L. Salcedo, Phys. Rev. D **69** (2004), 116003 [[arXiv:hep-ph/0312133](#)].

- 15) A. Vuorinen and L. G. Yaffe, Phys. Rev. D **74** (2006), 025011 [[arXiv:hep-ph/0604100](#)].
- 16) T. Banks and A. Ukawa, Nucl. Phys. B **225** (1983), 145.
- 17) J. Dai and V. P. Nair, Phys. Rev. D **74** (2006), 085014 [[arXiv:hep-ph/0605090](#)]; V. P. Nair, private communication.
- 18) O. Aharony, J. Marsano, S. Minwalla, K. Papadodimas, and M. Van Raamsdonk, Adv. Theor. Math. Phys. **8** (2004) 603 [[arXiv:hep-th/0310285](#)]; Phys. Rev. D **71** (2005), 125018 [[arXiv:hep-th/0502149](#)].